

Paper Reference(s)

**9801/01**

**Edexcel**

**Mathematics**

**Advanced Extension Award**

**Tuesday 25 June 2013 – Morning**

**Time: 3 hours**

**Materials required for examination**

Answer book (AB16)

Graph paper (ASG2)

Mathematical Formulae (Pink)

**Items included with question papers**

Nil

**Candidates may NOT use a calculator in answering this paper.**

**Instructions to Candidates**

In the boxes on the answer book provided, write the name of the examining body (Edexcel), your centre number, candidate number, the paper title (Mathematics), the paper reference (9801), your surname, initials and signature.

Check that you have the correct question paper.

Answers should be given in as simple a form as possible. e.g.  $\frac{2\pi}{3}$ ,  $\sqrt{6}$ ,  $3\sqrt{2}$ .

**Information for Candidates**

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 7 questions in this question paper.

The total mark for this paper is 100, of which 7 marks are for style, clarity and presentation.

There are 8 pages in this question paper. Any blank pages are indicated.

**Advice to Candidates**

You must ensure that your answers to parts of questions are clearly labelled.

You must show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

Printer's Log. No.

**P41807A**



*Turn over*

1. In the binomial expansion of

$$\left(1 + \frac{12n}{5}x\right)^n$$

the coefficients of  $x^2$  and  $x^3$  are equal and non-zero.

(a) Find the possible values of  $n$ .

(4)

(b) State, giving a reason, which value of  $n$  gives a valid expansion when  $x = \frac{1}{2}$

(2)

**(Total 6 marks)**

---

2. (a) Use the formula for  $\sin(A - B)$  to show that  $\sin(90^\circ - x) = \cos x$

(1)

(b) Solve for  $0 < \theta < 360^\circ$

$$2 \sin(\theta + 17^\circ) = \frac{\cos(\theta + 8^\circ)}{\cos(\theta + 17^\circ)}$$

(7)

**(Total 8 marks)**

---

3. The lines  $L_1$  and  $L_2$  have equations given by

$$L_1: \mathbf{r} = \begin{pmatrix} -7 \\ 7 \\ 1 \end{pmatrix} + \lambda \begin{pmatrix} 2 \\ 0 \\ -3 \end{pmatrix} \text{ and } L_2: \mathbf{r} = \begin{pmatrix} 7 \\ p \\ -6 \end{pmatrix} + \mu \begin{pmatrix} 10 \\ -4 \\ -1 \end{pmatrix}$$

where  $\lambda$  and  $\mu$  are parameters and  $p$  is a constant.

The two lines intersect at the point  $C$ .

(a) Find

(i) the value of  $p$ ,

(ii) the position vector of  $C$ .

(5)

(b) Show that the point  $B$  with position vector  $\begin{pmatrix} -13 \\ 11 \\ -4 \end{pmatrix}$  lies on  $L_2$ .

(1)

The point  $A$  with position vector  $\begin{pmatrix} -7 \\ 7 \\ 1 \end{pmatrix}$  lies on  $L_1$ .

(c) Find  $\cos(\angle ACB)$ , giving your answer as an exact fraction.

(3)

The line  $L_3$  bisects the angle  $ACB$ .

(d) Find a vector equation of  $L_3$ .

(4)

---

(Total 13 marks)

4. A sequence of positive integers  $a_1, a_2, a_3, \dots$  has  $r$ th term given by

$$a_r = 2^r - 1$$

(a) Write down the first 6 terms of this sequence. (1)

(b) Verify that  $a_{r+1} = 2a_r + 1$  (1)

(c) Find  $\sum_{r=1}^n a_r$  (3)

(d) Show that  $\frac{1}{a_{r+1}} < \frac{1}{2} \times \frac{1}{a_r}$  (1)

(e) Hence show that  $1 + \frac{1}{3} + \frac{1}{7} + \frac{1}{15} + \frac{1}{31} + \dots < 1 + \frac{1}{3} + \left(\frac{1}{7} + \frac{1}{7} + \frac{1}{7} + \dots\right)$  (2)

(f) Show that  $\frac{31}{21} < \sum_{r=1}^{\infty} \frac{1}{a_r} < \frac{34}{21}$  (5)

**(Total 13 marks)**

5. In this question  $u$  and  $v$  are functions of  $x$ . Given that  $\int u \, dx$ ,  $\int v \, dx$  and  $\int uv \, dx$  satisfy

$$\int uv \, dx = \left(\int u \, dx\right) \times \left(\int v \, dx\right) \quad uv \neq 0$$

(a) show that  $1 = \frac{\int u \, dx}{u} + \frac{\int v \, dx}{v}$  (3)

Given also that  $\frac{\int u \, dx}{u} = \sin^2 x$ ,

(b) use part (a) to write down an expression, in terms of  $x$ , for  $\frac{\int v \, dx}{v}$ , (1)

(c) show that 
$$\frac{1}{u} \frac{du}{dx} = \frac{1 - 2 \sin x \cos x}{\sin^2 x}$$
 (3)

(d) hence use integration to show that  $u = Ae^{-\cot x} \operatorname{cosec}^2 x$ , where  $A$  is an arbitrary constant. (6)

(e) By differentiating  $e^{\tan x}$  find a similar expression for  $v$ . (2)

**(Total 15 marks)**

6. (a) Starting from  $[f(x) - \lambda g(x)]^2 \geq 0$  show that  $\lambda$  satisfies the quadratic inequality

$$\left( \int_a^b [g(x)]^2 dx \right) \lambda^2 - 2 \left( \int_a^b f(x)g(x) dx \right) \lambda + \int_a^b [f(x)]^2 dx \geq 0$$

where  $a$  and  $b$  are constants and  $\lambda$  can take any real value.

(2)

- (b) Hence prove that

$$\left[ \int_a^b f(x)g(x) dx \right]^2 \leq \left[ \int_a^b [f(x)]^2 dx \right] \times \left[ \int_a^b [g(x)]^2 dx \right]$$

(3)

- (c) By letting  $f(x) = 1$  and  $g(x) = (1 + x^3)^{\frac{1}{2}}$  show that

$$\int_{-1}^2 (1 + x^3)^{\frac{1}{2}} dx \leq \frac{9}{2}$$

(4)

- (d) Show that  $\int_{-1}^2 x^2 (1 + x^3)^{\frac{1}{4}} dx = \frac{12\sqrt{3}}{5}$

(3)

- (e) Hence show that


$$\frac{144}{55} \leq \int_{-1}^2 (1 + x^3)^{\frac{1}{2}} dx$$

(4)

---

**(Total 16 marks)**

7.



**Figure 1**

Figure 1 shows a sketch of the curve  $C_1$  with equation  $y = f(x)$  where

$$f(x) = \frac{x}{3} + \frac{12}{x} \quad x \neq 0$$

The lines  $x = 0$  and  $y = \frac{x}{3}$  are asymptotes to  $C_1$ . The point  $A$  on  $C_1$  is a minimum and the point  $B$  on  $C_1$  is a maximum.

(a) Find the coordinates of  $A$  and  $B$ . (4)

There is a normal to  $C_1$ , which does not intersect  $C_1$  a second time, that has equation  $x = k$ , where  $k > 0$ .

(b) Write down the value of  $k$ . (1)

The point  $P(\alpha, \beta)$ ,  $\alpha > 0$  and  $\alpha \neq k$ , lies on  $C_1$ . The normal to  $C_1$  at  $P$  does not intersect  $C_1$  a second time.

(c) Find the value of  $\alpha$ , leaving your answer in simplified surd form. (5)

(d) Find the equation of this normal. (3)

The curve  $C_2$  has equation  $y = |f(x)|$

(e) Sketch  $C_2$  stating the coordinates of any turning points and the equations of any asymptotes. (4)

The line with equation  $y = mx + 1$  does not touch or intersect  $C_2$ .

(f) Find the set of possible values for  $m$ . (5)

**(Total 22 marks)**

**FOR STYLE, CLARITY AND PRESENTATION: 7 MARKS  
TOTAL FOR PAPER: 100 MARKS**

**END**

**BLANK PAGE**

**BLANK PAGE**