

Paper Reference(s)

**6666/01**

# **Edexcel GCE**

## **Core Mathematics C4**

### **Advanced Level**

**Monday 25 January 2010 – Morning**

**Time: 1 hour 30 minutes**

**Materials required for examination**

Mathematical Formulae (Pink or Green)

**Items included with question papers**

Nil

**Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation or integration, or have retrievable mathematical formulae stored in them.**

#### **Instructions to Candidates**

---

Write the name of the examining body (Edexcel), your centre number, candidate number, the unit title (Core Mathematics C4), the paper reference (6666), your surname, initials and signature.

#### **Information for Candidates**

---

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for the parts of questions are shown in round brackets, e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

#### **Advice to Candidates**

---

You must ensure that your answers to parts of questions are clearly labelled.

You must show sufficient working to make your methods clear to the Examiner.

Answers without working may not gain full credit.

1. (a) Find the binomial expansion of

$$\sqrt[3]{(1-8x)}, \quad |x| < \frac{1}{8},$$

in ascending powers of  $x$  up to and including the term in  $x^3$ , simplifying each term.

**(6)**

- (b) Show that, when  $x = \frac{1}{100}$ , the exact value of  $\sqrt[3]{(1-8x)}$  is  $\frac{\sqrt[3]{23}}{5}$ .


**(2)**

- (c) Substitute  $x = \frac{1}{100}$  into the binomial expansion in part (a) and hence obtain an approximation to  $\sqrt[3]{23}$ . Give your answer to 5 decimal places.

**(3)**

---

2.



**Figure 1**

Figure 1 shows a sketch of the curve with equation  $y = x \ln x$ ,  $x \geq 1$ . The finite region  $R$ , shown shaded in Figure 1, is bounded by the curve, the  $x$ -axis and the line  $x = 4$ .

The table shows corresponding values of  $x$  and  $y$  for  $y = x \ln x$ .

$x$	1	1.5	2	2.5	3	3.5	4
$y$	0	0.608			3.296	4.385	5.545

(a) Copy and complete the table with the values of  $y$  corresponding to  $x = 2$  and  $x = 2.5$ , giving your answers to 3 decimal places.

**(2)**

(b) Use the trapezium rule, with all the values of  $y$  in the completed table, to obtain an estimate for the area of  $R$ , giving your answer to 2 decimal places.

**(4)**

(c) (i) Use integration by parts to find  $\int x \ln x \, dx$ .

(ii) Hence find the exact area of  $R$ , giving your answer in the form  $\frac{1}{4}(a \ln 2 + b)$ , where  $a$  and  $b$  are integers.

**(7)**

3. The curve  $C$  has equation

$$\cos 2x + \cos 3y = 1, \quad -\frac{\pi}{4} \leq x \leq \frac{\pi}{4}, \quad 0 \leq y \leq \frac{\pi}{6}.$$

(a) Find  $\frac{dy}{dx}$  in terms of  $x$  and  $y$ .

**(3)**

The point  $P$  lies on  $C$  where  $x = \frac{\pi}{6}$ .

(b) Find the value of  $y$  at  $P$ .

**(3)**

(c) Find the equation of the tangent to  $C$  at  $P$ , giving your answer in the form  $ax + by + c\pi = 0$ , where  $a$ ,  $b$  and  $c$  are integers.

**(3)**

---

4. The line  $l_1$  has vector equation

$$\mathbf{r} = \begin{pmatrix} -6 \\ 4 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ -1 \\ 3 \end{pmatrix}$$

and the line  $l_2$  has vector equation

$$\mathbf{r} = \begin{pmatrix} -6 \\ 4 \\ -1 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ -4 \\ 1 \end{pmatrix}$$

where  $\lambda$  and  $\mu$  are parameters.

The lines  $l_1$  and  $l_2$  intersect at the point  $A$  and the acute angle between  $l_1$  and  $l_2$  is  $\theta$ .

- (a) Write down the coordinates of  $A$ .

**(1)**

- (b) Find the value of  $\cos \theta$ .

**(3)**

The point  $X$  lies on  $l_1$  where  $\lambda = 4$ .

- (c) Find the coordinates of  $X$ .

**(1)**

- (d) Find the vector  $\overrightarrow{AX}$ .

**(2)**

- (e) Hence, or otherwise, show that  $|\overrightarrow{AX}| = 4\sqrt{26}$ .

**(2)**

The point  $Y$  lies on  $l_2$ . Given that the vector  $\overrightarrow{YX}$  is perpendicular to  $l_1$ ,

- (f) find the length of  $AY$ , giving your answer to 3 significant figures.

**(3)**

---

5. (a) Find  $\int \frac{9x+6}{x} dx$ ,  $x > 0$ . (2)

(b) Given that  $y = 8$  at  $x = 1$ , solve the differential equation

$$\frac{dy}{dx} = \frac{(9x+6)y^{\frac{1}{3}}}{x}$$


giving your answer in the form  $y^2 = g(x)$ . (6)

---

6. The area  $A$  of a circle is increasing at a constant rate of  $1.5 \text{ cm}^2 \text{ s}^{-1}$ . Find, to 3 significant figures, the rate at which the radius  $r$  of the circle is increasing when the area of the circle is  $2 \text{ cm}^2$ . (5)

---

7.



**Figure 2**

Figure 2 shows a sketch of the curve  $C$  with parametric equations

$$x = 5t^2 - 4, \quad y = t(9 - t^2)$$

The curve  $C$  cuts the  $x$ -axis at the points  $A$  and  $B$ .

(a) Find the  $x$ -coordinate at the point  $A$  and the  $x$ -coordinate at the point  $B$ . **(3)**

The region  $R$ , as shown shaded in Figure 2, is enclosed by the loop of the curve.


(b) Use integration to find the area of  $R$ . **(6)**

---

8. (a) Using the substitution  $x = 2 \cos u$ , or otherwise, find the exact value of

$$\int_1^{\sqrt{2}} \frac{1}{x^2 \sqrt{4-x^2}} dx.$$

(7)



**Figure 3**

Figure 3 shows a sketch of part of the curve with equation  $y = \frac{4}{x(4-x^2)^{\frac{1}{4}}}$ ,  $0 < x < 2$ .

The shaded region  $S$ , shown in Figure 3, is bounded by the curve, the  $x$ -axis and the lines with equations  $x = 1$  and  $x = \sqrt{2}$ . The shaded region  $S$  is rotated through  $2\pi$  radians about the  $x$ -axis to form a solid of revolution.

- (b) Using your answer to part (a), find the exact volume of the solid of revolution formed.

(3)

---

**TOTAL FOR PAPER: 75 MARKS**

**END**




**January 2010**  
**6666 Core Mathematics C4**  
**Mark Scheme**

Question Number	Scheme	Marks
Q1	$(a) (1-8x)^{\frac{1}{2}} = 1 + \left(\frac{1}{2}\right)(-8x) + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)}{2}(-8x)^2 + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)}{3!}(-8x)^3 + \dots$ $= 1 - 4x - 8x^2; -32x^3 - \dots$	M1 A1 A1; A1      (4)
	$(b) \sqrt{(1-8x)} = \sqrt{\left(1 - \frac{8}{100}\right)}$ $= \sqrt{\frac{92}{100}} = \sqrt{\frac{23}{25}} = \frac{\sqrt{23}}{5} \quad *$	M1 cs0 A1      (2)
	$(c) 1 - 4x - 8x^2 - 32x^3 = 1 - 4(0.01) - 8(0.01)^2 - 32(0.01)^3$ $= 1 - 0.04 - 0.0008 - 0.00032 = 0.959168$ $\sqrt{23} = 5 \times 0.959168$ $= 4.79584$	M1 M1 cao A1      (3) <b>[9]</b>

Question Number	Scheme	Marks
Q2	(a) 1.386, 2.291 <span style="float: right;">awrt 1.386, 2.291</span>	B1 B1 (2)
	(b) $A \approx \frac{1}{2} \times 0.5( \dots )$ $= \dots (0 + 2(0.608 + 1.386 + 2.291 + 3.296 + 4.385) + 5.545)$ $= 0.25(0 + 2(0.608 + 1.386 + 2.291 + 3.296 + 4.385) + 5.545)$ ft their (a) $= 0.25 \times 29.477 \dots \approx 7.37$	B1 M1 A1ft A1 (4)
	(c)(i) $\int x \ln x \, dx = \frac{x^2}{2} \ln x - \int \frac{x^2}{2} \times \frac{1}{x} \, dx$ $= \frac{x^2}{2} \ln x - \int \frac{x}{2} \, dx$ $= \frac{x^2}{2} \ln x - \frac{x^2}{4} (+C)$	M1 A1 M1 A1
	(ii) $\left[ \frac{x^2}{2} \ln x - \frac{x^2}{4} \right]_1^4 = (8 \ln 4 - 4) - \left( -\frac{1}{4} \right)$ $= 8 \ln 4 - \frac{15}{4}$ $= 8(2 \ln 2) - \frac{15}{4}$ <span style="margin-left: 2em;"><math>\ln 4 = 2 \ln 2</math> seen or implied</span> $= \frac{1}{4}(64 \ln 2 - 15)$ <span style="margin-left: 2em;"><math>a = 64, b = -15</math></span>	M1 M1 A1 (7)
		<b>[13]</b>

Question Number	Scheme	Marks
Q3	(a) $-2 \sin 2x - 3 \sin 3y \frac{dy}{dx} = 0$ $\frac{dy}{dx} = -\frac{2 \sin 2x}{3 \sin 3y}$ Accept $\frac{2 \sin 2x}{-3 \sin 3y}, \frac{-2 \sin 2x}{3 \sin 3y}$	M1 A1 A1      (3)
	(b) At $x = \frac{\pi}{6}$ , $\cos\left(\frac{2\pi}{6}\right) + \cos 3y = 1$ $\cos 3y = \frac{1}{2}$ $3y = \frac{\pi}{3} \Rightarrow y = \frac{\pi}{9}$ awrt 0.349	M1 A1 A1      (3)
	(c) At $\left(\frac{\pi}{6}, \frac{\pi}{9}\right)$ , $\frac{dy}{dx} = -\frac{2 \sin 2\left(\frac{\pi}{6}\right)}{3 \sin 3\left(\frac{\pi}{9}\right)} = -\frac{2 \sin \frac{\pi}{3}}{3 \sin \frac{\pi}{3}} = -\frac{2}{3}$ $y - \frac{\pi}{9} = -\frac{2}{3}\left(x - \frac{\pi}{6}\right)$ Leading to $6x + 9y - 2\pi = 0$	M1 M1 A1      (3) [9]

Question Number	Scheme	Marks
Q4	(a) $A: (-6, 4, -1)$ <span style="float: right;">Accept vector forms</span>	B1 (1)
	(b) $\begin{pmatrix} 4 \\ -1 \\ 3 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ -4 \\ 1 \end{pmatrix} = 12 + 4 + 3 = \sqrt{4^2 + (-1)^2 + 3^2} \sqrt{3^2 + (-4)^2 + 1^2} \cos \theta$ $\cos \theta = \frac{19}{26}$ <span style="float: right;">awrt 0.73</span>	M1 A1 A1 (3)
	(c) $X: (10, 0, 11)$ <span style="float: right;">Accept vector forms</span>	B1 (1)
	(d) $\vec{AX} = \begin{pmatrix} 10 \\ 0 \\ 11 \end{pmatrix} - \begin{pmatrix} -6 \\ 4 \\ -1 \end{pmatrix}$ $= \begin{pmatrix} 16 \\ -4 \\ 12 \end{pmatrix}$ <span style="float: right;">Either order cao</span>	M1 A1 (2)
	(e) $ \vec{AX}  = \sqrt{16^2 + (-4)^2 + 12^2}$ $= \sqrt{416} = \sqrt{16 \times 26} = 4\sqrt{26} *$ <span style="float: right;">Do not penalise if consistent incorrect signs in (d)</span>	M1 A1 (2)
	(f)  <span style="float: right;">Use of correct right angled triangle</span> $\frac{ \vec{AX} }{d} = \cos \theta$ $d = \frac{4\sqrt{26}}{\frac{19}{26}} \approx 27.9$ <span style="float: right;">awrt 27.9</span>	M1 M1 A1 (3) <b>[12]</b>

Question Number	Scheme	Marks
Q5	(a) $\int \frac{9x+6}{x} dx = \int \left(9 + \frac{6}{x}\right) dx$ $= 9x + 6 \ln x (+C)$	M1 A1 (2)
	(b) $\int \frac{1}{y^{\frac{1}{3}}} dy = \int \frac{9x+6}{x} dx$ $\int y^{-\frac{1}{3}} dy = \int \frac{9x+6}{x} dx$	Integral signs not necessary B1
	$\frac{y^{\frac{2}{3}}}{\frac{2}{3}} = 9x + 6 \ln x (+C)$	$\pm ky^{\frac{2}{3}} =$ their (a) M1
	$\frac{3}{2} y^{\frac{2}{3}} = 9x + 6 \ln x (+C)$	ft their (a) A1ft
	$y = 8, x = 1$ $\frac{3}{2} 8^{\frac{2}{3}} = 9 + 6 \ln 1 + C$	M1
	$C = -3$ $y^{\frac{2}{3}} = \frac{2}{3}(9x + 6 \ln x - 3)$ $y^2 = (6x + 4 \ln x - 2)^3 \quad (= 8(3x + 2 \ln x - 1)^3)$	A1 A1 (6) [8]

Question Number	Scheme	Marks
Q6	$\frac{dA}{dt} = 1.5$ $A = \pi r^2 \Rightarrow \frac{dA}{dr} = 2\pi r$ <p>When <math>A = 2</math></p> $2 = \pi r^2 \Rightarrow r = \sqrt{\frac{2}{\pi}} (= 0.797\ 884 \dots)$ $\frac{dA}{dt} = \frac{dA}{dr} \times \frac{dr}{dt}$ $1.5 = 2\pi r \frac{dr}{dt}$ $\frac{dr}{dt} = \frac{1.5}{2\pi\sqrt{\frac{2}{\pi}}} \approx 0.299$	<p>B1</p> <p>B1</p> <p>M1</p> <p>M1</p> <p>A1</p> <p style="text-align: right;">awrt 0.299</p> <p style="text-align: right;"><b>[5]</b></p>

Question Number	Scheme	Marks
Q7	<p>(a) <math>y = 0 \Rightarrow t(9 - t^2) = t(3 - t)(3 + t) = 0</math>  <math>t = 0, 3, -3</math></p> <p>At <math>t = 0</math>, <math>x = 5(0)^2 - 4 = -4</math>  At <math>t = 3</math>, <math>x = 5(3)^2 - 4 = 41</math>  (At <math>t = -3</math>, <math>x = 5(-3)^2 - 4 = 41</math>)  At <math>A</math>, <math>x = -4</math>; at <math>B</math>, <math>x = 41</math></p> <p>(b) <math>\frac{dx}{dt} = 10t</math></p> $\int y \, dx = \int y \frac{dx}{dt} \, dt = \int t(9 - t^2)10t \, dt$ $= \int (90t^2 - 10t^4) \, dt$ $= \frac{90t^3}{3} - \frac{10t^5}{5} (+C) \quad (= 30t^3 - 2t^5 (+C))$ $\left[ \frac{90t^3}{3} - \frac{10t^5}{5} \right]_0^3 = 30 \times 3^3 - 2 \times 3^5 \quad (= 324)$ $A = 2 \int y \, dx = 648 \quad (\text{units}^2)$	<p>Any one correct value B1  Method for finding one value of x M1</p> <p>Both A1 (3)</p> <p>Seen or implied B1</p> <p>M1 A1</p> <p>A1</p> <p>M1</p> <p>A1 (6)  <b>[9]</b></p>

Question Number	Scheme	Marks
Q8	<p>(a) <math>\frac{dx}{du} = -2 \sin u</math></p> $\int \frac{1}{x^2 \sqrt{4-x^2}} dx = \int \frac{1}{(2 \cos u)^2 \sqrt{4-(2 \cos u)^2}} \times -2 \sin u du$ $= \int \frac{-2 \sin u}{4 \cos^2 u \sqrt{4 \sin^2 u}} du \quad \text{Use of } 1 - \cos^2 u = \sin^2 u$ $= -\frac{1}{4} \int \frac{1}{\cos^2 u} du \quad \pm k \int \frac{1}{\cos^2 u} du$ $= -\frac{1}{4} \tan u (+C) \quad \pm k \tan u$ <p><math>x = \sqrt{2} \Rightarrow \sqrt{2} = 2 \cos u \Rightarrow u = \frac{\pi}{4}</math></p> <p><math>x = 1 \Rightarrow 1 = 2 \cos u \Rightarrow u = \frac{\pi}{3}</math></p> $\left[ -\frac{1}{4} \tan u \right]_{\frac{\pi}{3}}^{\frac{\pi}{4}} = -\frac{1}{4} \left( \tan \frac{\pi}{4} - \tan \frac{\pi}{3} \right)$ $= -\frac{1}{4} (1 - \sqrt{3}) \quad \left( = \frac{\sqrt{3}-1}{4} \right)$ <p>(b) <math>V = \pi \int_1^{\sqrt{2}} \left( \frac{4}{x(4-x^2)^{\frac{1}{4}}} \right)^2 dx</math></p> $= 16\pi \int_1^{\sqrt{2}} \frac{1}{x^2 \sqrt{4-x^2}} dx \quad 16\pi \times \text{integral in (a)}$ $= 16\pi \left( \frac{\sqrt{3}-1}{4} \right) \quad 16\pi \times \text{their answer to part (a)}$	<p>B1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>M1</p> <p>A1 (7)</p> <p>M1</p> <p>M1</p> <p>A1ft (3)</p> <p>[10]</p>